arXiv:1701.05643v1 [math.NT] 19 Jan 2017

# A note on primitive 1−normal elements over finite fields

Lucas Reis

*Departamento de Matemática, Universidade Federal de Minas Gerais, UFMG, Belo Horizonte, MG, 30123-970, Brazil*

**Abstract**

Let $q$ be a prime power of a prime $p$, $n$ a positive integer and $\mathbb{F}_{q^n}$ the finite field with $q^n$ elements. The $k$−normal elements over finite fields were introduced and characterized by Huczynska *et al* (2013). Under the condition that $n$ is not divisible by $p$, they obtained an existence result on primitive 1−normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ for $q > 2$. In this note, we extend their result to the excluded case $q = 2$.

*Keywords:* Finite Fields, Normal Basis, k-normal elements, Primitive elements
2010 MSC: 12E20, 11T06

## 1. Introduction

Let $\mathbb{F}_{q^n}$ be the finite field with $q^n$ elements, where $q$ is a prime power and $n$ is a positive integer. Recall that an element $\alpha \in \mathbb{F}_{q^n}$ is said to be normal over $\mathbb{F}_q$ if $A = \{\alpha, \alpha^q, \cdots, \alpha^{q^{n-1}}\}$ is a basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$; A is called a normal basis. Normal basis are frequently used in cryptography and computer algebra systems; sometimes it is useful to take normal basis composed by primitive elements, i.e., generators of the multiplicative group $\mathbb{F}_{q^n}^*$. The *Primitive Normal Basis Theorem* states that for any extension field $\mathbb{F}_{q^n}$ of $\mathbb{F}_q$, there exists a basis composed by primitive normal elements; this result was first proved by Lenstra and Schoof [6] and a proof without the use of a computer was latter given in [3].

A characterization of normal elements is given in ([7], Theorem 2.39): an element $\alpha \in \mathbb{F}_{q^n}$ is normal over $\mathbb{F}_q$ if and only if the polynomials

$$g_\alpha(x) := \sum_{i=0}^{n-1} \alpha^{q^i} x^{n-1-i} \quad \text{and} \quad x^n - 1,$$

are relatively prime over $\mathbb{F}_{q^n}$. Motivated by this characterization, in [5], the authors introduce $k$−normal elements:

*Email address:* lucasreismat@gmail.com (Lucas Reis)

**Definition 1.1.** *Let $\alpha \in \mathbb{F}_{q^n}^*$ and $g_\alpha(x) = \sum_{i=0}^{n-1} \alpha^{q^i} \cdot x^{n-1-i}$. We say that $\alpha$ is $k-$normal over $\mathbb{F}_q$ if the greatest common divisor of $x^n - 1$ and $g_\alpha(x)$ over $\mathbb{F}_{q^n}$ has degree $k$.*

From definition, $0-$normal elements correspond to normal elements in the usual sense. In the same paper, the authors give a characterization of $k-$normal elements and find a formula for their number. Also, they obtain an existence result on primtive $1-$normal elements:

**Theorem 1.2** ([5], Theorem 5.10)**.** *Let $q = p^e$ be a prime power and $n$ a positive integer not divisible by $p$. Assume that $n \geq 6$ if $q \geq 11$ and that $n \geq 3$ if $3 \leq q \leq 9$. Then there exists a primitive $1-$normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.*

The authors propose an extension of the above theorem for all pairs $(q, n)$ with $n \geq 2$ as a problem ([5], Problem 6.2); they conjectured that such elements always exist. However, it was proved in [1] that for odd $q > 3$ and $n = 2$, there are no primitive $1-$normal elements of $\mathbb{F}_{q^2}$ over $\mathbb{F}_q$. The aim of this note is to extend Theorem 1.2 to the case when $q = 2$ and $n$ is odd. Essentially, we show that the tools used in [5] to prove Theorem 1.2 can be adapted to that case.

## 2. Existence of primitive $1-$normal elements over $\mathbb{F}_2$

First, we present some definitions and results that will be useful in the rest of this paper.

**Definition 2.1.**

(a) *Let $f(x)$ be a monic polynomial with coefficients in $\mathbb{F}_q$. The Euler Phi Function for polynomials over $\mathbb{F}_q$ is given by*

$$\Phi_q(f) = \left| \left( \frac{\mathbb{F}_q[x]}{\langle f \rangle} \right)^* \right|,$$

*where $\langle f \rangle$ is the ideal generated by $f(x)$ in $\mathbb{F}_q[x]$.*

(b) *If $t$ is a positive integer (or a monic polynomial over $\mathbb{F}_q$), $W(t)$ denotes the number of square-free (monic) divisors of $t$.*

We have an interesting formula for the number of $k-$normal elements over finite fields:

**Lemma 2.2** ([5], Theorem 3.5)**.** *The number $N_k$ of $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is given by*

$$\sum_{\substack{h | x^n - 1 \\ \deg(h) = n-k}} \Phi_q(h), \tag{1}$$

*where the divisors are monic and polynomial division is over $\mathbb{F}_q$.*

In particular, if $n \geq 2$, the number of $1-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is at least equal to $\Phi_q(T)$, where $T = \frac{x^n-1}{x-1}$.

*2.1. A sieve inequality*

The proof of Theorem 1.2 is based in an application of the *Lenstra-Schoof* method, introduced in [6]; this method has been used frequently in the characterization of elements in finite fields with particular properties like being primitive, normal and of zero-trace. For more details, see [3] and [5]. In particular, from Corollary 5.8 of [5], we can easily deduce the following:

**Lemma 2.3.** *Suppose that $q$ is a power of a prime $p$, $n \geq 2$ is a positive integer not divisible by $p$ and $T(x) = \frac{x^n-1}{x-1}$. If*

$$W(T) \cdot W(q^n - 1) < q^{n/2-1}, \tag{2}$$

*then there exist $1-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.*

Inequality (2) is an essential step in the proof of Theorem 1.2 and it was first studied in [2]; under the condition that $n \geq 6$ for $q \geq 11$ and $n \geq 3$ for $3 \leq q \leq 9$, this inequality is not true only for a finite number of pairs $(q, n)$ (see Theorem 4.5 of [2]). Here we extend the study of inequality (2) to the case when $q = 2$ and $n$ is odd. First, we have the following:

**Proposition 2.4.** *Suppose that $n \geq 3$ is odd and $T(x) := \frac{x^n-1}{x-1} \in \mathbb{F}_2[x]$. Then $W(T) \leq 2^{\frac{n+9}{5}}$.*

*Proof.* For each $2 \leq i \leq 4$, let $s_i$ be the number of irreducible factors of degree $i$ dividing $T(x)$. Since $n$ is odd, $T(x)$ has no linear factor. By a direct verification we see that the number of irreducible polynomials over $\mathbb{F}_2$ of degrees $2, 3$ and $4$ is $1, 2$ and $3$, respectively. Hence $s_2 \leq 1$, $s_3 \leq 2$ and $s_4 \leq 3$. In particular, the number of irreducible factors of $T(x)$ over $\mathbb{F}_2$ is at most

$$\frac{n - 1 - 2s_2 - 3s_3 - 4s_4}{5} + s_2 + s_3 + s_4 = \frac{n - 1 + 3s_2 + 2s_3 + s_4}{5}.$$

Since $\frac{n-1+3s_2+2s_3+s_4}{5} \leq \frac{n-1+3+4+3}{5} = \frac{n+9}{5}$, we conclude the proof. □

According to Lemma 7.5 in [4], $W(2^n - 1) < 2^{\frac{n}{7}+2}$ if $n$ is odd. In particular, we obtain the following:

**Corollary 2.5.** *Suppose that $n \neq 15$ is odd, $q = 2$ and $T(x) = \frac{x^n-1}{x-1} \in \mathbb{F}_2[x]$. For $n > 9$, inequality* (2) *holds.*

*Proof.* Notice that $\frac{n+9}{5}+\frac{n}{7}+2<\frac{n}{2}-1$ for $n\geq 31$. From Proposition 2.4 and Lemma 7.5 of [4], it follows that inequality (2) holds for odd $n\geq 31$. The remaining cases can be verified directly. □

We are ready to state and prove our result:

**Theorem 2.6.** *Suppose that $n\geq 3$ is odd. Then there exist a primitive $1-$normal element of $\mathbb{F}_{2^n}$ over $\mathbb{F}_2$.*

*Proof.* According to Lemma 2.3 and Corollary 2.5, this statement is true for $n>9$ if $n\neq 15$. For the remaining cases $n=3,5,7,9$ and 15 we use the following argument. Let $P$ be the number of primitive elements of $\mathbb{F}_{2^n}$ and $N_1$ the number of $1-$normal elements of $\mathbb{F}_{2^n}$ over $\mathbb{F}_2$; if $P+N_1>2^n$, there exists a primitive $1-$normal element of $\mathbb{F}_{2^n}$ over $\mathbb{F}_2$. Notice that $P=\varphi(2^n-1)$ and, according to Lemma 2.2, $N_1\geq\Phi_2\left(\frac{x^n-1}{x-1}\right)$. By a direct calculation we see that

$$\varphi(2^n-1)+\Phi_2\left(\frac{x^n-1}{x-1}\right)>2^n,$$

for $n=3,5,7,9$ and 15. This completes the proof. □

## References


[1] M. Alizadeh. *Some notes on the $k-$normal elements and $k-$normal polynomials over finite fields*, Journal of Algebra and Its Applications 16 (2017).

[2] S. D. Cohen, D. Hachenberger, *Primitive normal bases with prescribed trace*, Applicable Algebra in Engineering, Communication and Computing 9 (1999) 383403.

[3] S. D. Cohen, S. Huczynska. *The primitive normal basis theorem - without a computer*, Journal of the London Mathematical Society 67 (2003) 41-56.

[4] S. D. Cohen. *Pairs of primitive elements in fields of even order*, Finite Fields Appl. 28 (2014) 22-42.

[5] S. Huczynska, G.L. Mullen, D. Panario, and D. Thomson, *Existence and properties of $k-$normal elements over finite fields*, Finite Fields Appl. 24 (2013) 170-183.

[6] H. W. Lenstra, R. Schoof, *Primitive normal bases for finite fields*, Mathematics of Computation 48 (1987) 217-231.

[7] R. Lidl, H. Niederreiter, *Finite Fields: Encyclopedia of Mathematics and Its Applications*, vol. 20, 2nd ed. Cambridge University Pres, Cambridge, 1997.